Acta Aquatica Turcica
E-ISSN: 2651-5474 17(1), 108-118 (2021) DOI:https://doi.org/10.22392/actaquatr.762038

Effect of Mediterranean Mussels *(Mytilus galloprovincialis)* From Polluted Areas on Hepatotoicity in Rats by Immunuhistochemical Method

Latife Ceyda İRKİN[1], Suat ÇAKINA[2*], İlhan ÖZDEMİR[3], Şamil ÖZTÜRK[2]

[1]Çanakkale Onsekiz Mart University, Çanakkale Faculty of Applied Sciences, Department of Fisheries Technology, Çanakkale, Turkey,
[2*]Çanakkale Onsekiz Mart University, Vocational School of Health Services, Çanakkale, Turkey,
[3]Erzurum Atatürk University, Faculty of Medicine, Department of Obstetrics and Gynecology, Erzurum, Turkey,
[2]Çanakkale Onsekiz Mart University, Vocational School of Health Services, Çanakkale, Turkey,

***Corresponding Author:** suatcakina@comu.edu.tr

Research Article

Received 01 July 2020; Accepted 22 October 2020; Release date 01 March 2021.

How to Cite: İrkin, L. C., Çakına, S., Özdemir, İ., & Öztürk, Ş. (2021). Effect of Mediterranean Mussels *(Mytilus galloprovincialis)* from polluted areas on hepatotoicity in rats by immunuhistochemical method. *Acta Aquatica Turcica, 17*(1), 108-118. https://doi.org/10.22392/actaquatr.762038

Abstract

Mussels *(Mytilus galloprovincialis*) are aquatic organisms that can accumulate all the factors that pollute the fresh and saltwater environment. Although the Dardanelles has been exposed to environmental pollution for many years, it is also an important region for crustaceans. We aimed to investigate the histopathological changes in the liver parenchyma which is an important gland of the digestive system by feeding these mussels to rats. Twenty-four male Wistar albino were used in the study. The first group (control): fed with standard rat feed, the second group (experiment 1): 4/5 mussel + 1/5 standard rat feed daily; third group (experiment 2): 4/5 mussel + 1/5 standard rat feed every other day, the fourth group (experiment 3): groups were formed with 4/5 mussel + 1/5 standard rat feed every three days. All liver tissue samples taken from the experimental and control groups were stained with hematoxylin-eosin, and immuohitochemically staining inflammatory marker TNF-α and NF-κB after routine histopathological follow-up and analyzed with a light microscope image analysis system. It was observed that mononuclear cells caused inflammation of portal areas, increased sinusoidal dilatation and congestion and degeneration due to vacuolization in hepatocytes in the liver parenchyma of mussel-fed rats. Besides, immunohistochemical staining, TNF-α, and NF-κB immunoreactivity were observed in the liver cells of especially in the second group of rats. As a result, it has been shown that the consumption of mussels obtained and marketed without considering environmental pollution may trigger important digestive system organs of liver diseases.

Keywords: Immunohistochemistry, inflammation, liver, hepatocyte degeneration, mussels.

Kirlenmiş Alanlardan Toplanan Akdeniz Midyesi'nin Ratlarda İmmünohistokimyasal Metotla Hepatotoksik Etkisi

Özet

Midye (*Mytilus galloprovincialis*), tatlı ve tuzlu su ortamını kirleten tüm faktörleri biriktirebilen sucul organizmalardır. Çanakkale Boğazı uzun yıllar boyunca çevre kirliliğine maruz kalsada, kabuklular için de önemli bir bölgedir. Bu midyelerin sindirim sisteminin önemli bir bezi olan karaciğer parankimindeki histopatolojik etkilerini araştırmayı amaçladık. Çalışmada 24 adet erkek Wistar albino türü rat kullanıldı. Birinci grup (kontrol): standart sıçan yemi ile beslendi, ikinci grup (deney 1): hergün 4/5 midye + 1/5 standart sıçan yemi; üçüncü grup (deney 2): 4/5 midye + 1/5 standart sıçan yemi 2 günde bir, dördüncü grup (deney 3): gruplar 4/5 midye + 1/5 standart sıçan yemi ile her üç günde bir oluşturuldu. Deney ve kontrol gruplarından alınan tüm karaciğer doku örnekleri, histopatolojik takipten sonra rutin hematoksilin-eozin ve enflamatuar belirteç olan TNF-α ve NF-κB ile immünohistokimyasal boyama yapıldı ve ışık mikroskobu görüntü analiz sistemi ile analiz edildi. Mononükleer hücrelerin portal alanlarda yangıya sebep olduğu, karaciğer parankimindeki hepatositlerde vakuolizasyonun dejenerasyon sonucu meydana geldiği, santral ven ve sinüzoidal dilatasyon ve konjesyonun olduğu gözlemlendi. Ek olarak, özellikle ikinci sıçan grubunun karaciğer hücrelerinde immünohistokimyasal boyama, TNF-α ve NF-κB şiddetli immünoreaktivitesi gözlendi. Sonuç olarak, çevre kirliliği dikkate alınmadan elde edilen ve pazarlanan midye tüketiminin sindirim sisteminin önemli bir organı olan karaciğer hastalıklarını tetikleyebileceği gösterilmiştir.

Anahtar kelimeler: İmmünohistokimya, inflamasyon, karaciğer, hepatosit hasarı, midye

INTRODUCTION

The relationship between healthy food and quality of life is gaining importance day by day. Accordingly, it increases consumers' orientation towards natural and delicious products that are not contaminated with drugs or chemicals. Despite all these positive thoughts, the increasing population every day and the increase in food consumption parallel to this increase in the orientation towards alternative food sources. Consumption in fish and shellfish occasionally bursts. Especially in coastal areas, there are seashells that have become food culture. Creatures such as mussels, oysters, and sea urchins are among the more preferred products as nutrients. Mussels are organisms that feed on organic and inorganic substances in the water. Mussels can also filter toxic substances during water filtration. Because of these properties, they can store harmful substances in water in their bodies al. (Figueras et al., 2019). The mussels are fed with microorganisms that secrete neurotoxins, especially in the summer months. Also, they filter the water and accumulate heavy metals, pesticides, and bacteria in the tissues. As a result of the pollution of seawater with environmental factors, toxin levels increase in the body of these creatures (Voudanta et al., 2016). Dardanelles is the marine area where the mussel population is the most and it is very prone to contamination in terms of location. The seas have indeed been polluted for years with domestic waste, pesticides, and industrial waste (Yigit et al., 2018).

Most of the liver studies have been conducted to examine regeneration as a result of either experimental or chemical damage (Palmes & Spiegel, 2004). TNF-α plays a role in the pathophysiology of TNF-α, viral hepatitis, alcoholic liver disease, non-alcoholic fatty liver disease and, ischemia-reperfusion injury in the liver. Studies have shown that when the liver enters the degenerative process, TNF-α induces increased synthesis in hepatocytes and Kupffer cells in the cytoplasms (Yang & Seki, 2015) When oxidative stress in tissues and associated damage to cells increases, activation of the TNF-α receptor is followed by activation of the nuclear factor kappa B (NF-κB). Thus, NF-κB goes to the nucleus, and then NF-κB activates genes that try to block TNF-induced apoptosis. In resting cells, NF-κB is an inactive form in the cytoplasm (Karin, 2006; Schattenberg & Schuchmann, 2009).

Heavy metals are defined as metallic elements with a relatively high density than water (Fergusson, 1990). Heavy metals contain metalloids and heavy metals such as arsenic can have toxic effects even at low doses (Pourret & Hursthouse, 2019). In recent years, there has been an increasing ecological and global public health concern associated with environmental pollution by these metals. Besides, human exposure has increased significantly as their use has increased exponentially in some industrial, agricultural, domestic and technological applications (Tchounwou et al., 2012). In biological systems, it has been reported that heavy metals affect cellular organelles and components such as cell membranes, mitochondria, lysosomes, endoplasmic reticulum, nuclei, and enzymes involved in metabolism, detoxification, and damage repair (Wang & Shi, 2001). Metal ions have been found to interact with cell components such as DNA and nuclear proteins and cause conformational changes that can lead to DNA damage and cell cycle modulation, carcinogenesis, or apoptosis (Chang et al., 1996; Hubbard, 2005). Various forms of heavy metals, pesticides, viral and bacterial organisms that crustaceans filter and accumulate in their tissues also pass to mammals that consume these creatures (Gorinstein et al., 2008). As a result, it will inevitably produce important histopathological results in the liver. In this study, we aimed to draw attention to the consumption of seafood consumption in a healthier and more reliable environment by showing the changes in the liver tissue with histochemistry and immunohistochemical techniques by giving them to the rats with an experimental nutrition model in vivo.

MATREIALS and METHODS

The mussels used in the study were removed by diving from various locations determined in the Dardanelles of April- May 2019. Mussels were selected from the same region and those close to the same size. Mussel muscle tissue was assumed to be contaminated based on analysis results. The mussels were boiled in the shell and after the water was taken in the oven, they were turned into o pellet and fed to the subjects as feed. In this study, 24 male Wistar albino rats (250-300 g in weight) were used. All rats were housed in a 12-hour light and 12-hour dark environment with an average temperature of 22 ± 1°C, humidity 55 ± 5, ventilation and air conditioning system. Rats were given as

much water as they could drink. Standard rat food and mussel were given according to 15% of the weight of each rat in feeding planning (Gezen, 2018).

Experimental groups

The first group (Control, n = 6); Standard rat food,

The second group (Experiment 1, n = 6); 4/5 mussels + 1/5 standard rat food daily,

The third group (Experiment 2, n = 6); 4/5 mussels + 1/5 standard rat food every two days; other day standard rat food,

The fourth group (Experiment 3, n = 6); 4/5 mussels + 1/5 standard rat food every three days; the other two days are standard rat food; It was fed for four weeks.

Ethics Statement

A total of 24 male Wistar albino rats were used in the study. The study protocol was approved by the Canakkale Onsekiz Mart University Ethics Committee for Animal Research (Protocol number: 2020/04-07).

Histopathological Examination

30 days after the start of the study, rats were removed for this study after rats were sacrificed under ketas (150 mg/kg; i.p) and alfazyne 2% (25 mg/kg; i.p) anesthesia. Liver tissue samples from all groups were detected in a 10% neutral buffered formaldehyde solution (Bio Optica) for 24 hours. Tissue samples were passed through graded alcohols and their juices were removed. Then, the tissues passed through xylene were made transparent and the alcohols of the tissues were removed. Paraffin was allowed to enter into the tissue samples passed through xylene + paraffin and paraffin stages in a 60°C oven. Tissue samples removed from paraffin were blocked using a tissue embedding device. Tissue samples taken to the blocks were cut 4-5 microns thick in microtome for routine histopathological staining and taken into a water bath. The tissue samples opened here were taken on a normal slide and Hematoxylin-Eosin staining was applied.

Immunohistochemically Examination

Immunohistochemically reactions were performed according to the ABC technique described. Following this step, the sections were incubated with a polyclonal nuclear factor Kappa-B (NF-κB p50, Abcam), tumor necrosis factor (TNF-α, Abcam), then the sections were incubated with biotinylated anti-mouse Immunoglobulin-G (DAKO LSAB 2 Kit, Invitrogen). Following this step, the sections were incubated with the ABC complex (DAKO LSAB 2 Kit). For background staining, Mayer's Hematoxylin (Gürpınar et al., 2012; Öztürk et al., 2019).

Evaluation of tissue samples and statistics

During the evaluation of the results, the immunoreactivity was evaluated with the H-score method, calculating the ratio of immunopositivity cells to all cells in the selected fields. Immunoreactive cell count was performed by a blinded observer and graded as follows: 0 denoted no staining; 1 denoted weakly; 2 denoted moderate; 3 denoted strong staining in a specified field. The respective score was then calculated using the following formula: H-score = (% stained cells at 0) x 0 + (%stained cells at 1+) x 1 + (%stained cells at 2+) x 2 + (%stained cells at 3+) x3. The H-score value varies from 0 to 300. SPSS 19 version applied for statistical evaluation of the results obtained with this formula. To determine the differences NF-κB and TNF-α immunoreactivities between groups, the Kruskal –Wallis Test, which is one of the nonparametric tests, will be used. $P<0.05$ Difference between the groups will be considered significant.

RESULT and DISCUSSION

Histophatological Findings

Control Group: Hepatocellular damage, biliary tract damage, vascular damage, and sinusoid cell damage, and no histopathological findings of tumor cases were observed when staining liver tissue samples from rats fed with standard rat food every day with Hematoxylin and Eosin (Figure 1). Remark cord structure, sinusoid, and central vein, and portal vein, portal artery, and bile ducts located in the portal area were observed to have normal histological structure.

Experimental groups: The following histopathology table occurred in all experimental groups, respectively.

a- Hepatocellular Damage; Vacuolar degeneration, focal necrosis, inflammatory cells including neutrophils and eosinophils in the lobule and portal area were detected. Vacuolar degeneration in most of the hepatocytes manifested by diffuse swelling, pale staining of the cytoplasm, and the appearance

of the cytoplasmic residues around the nucleus. In some hepatocytes, the picnotic nucleus was observed, lobular inflammation in small foci, periductal inflammation with portal inflammation (Figure 1, Table 1).

Table 1. Histopathological evaluation of liver tissue samples.

	Groups			
Parameters	**Control**	**Experiment 1 (Every day giving mussel)**	**Experiment 2 (Two day giving mussel)**	**Experiment 3 (Three day giving mussel)**
Congestion	-	++++	++	+
Dilatation	+	++++	+++	++
Inflammation	-	+++	++	+

b- Vascular Damage; Central, portal, and sinusoidal dilatation, congestion in portal veins, central veins and sinusoid were observed. These histophatological changes occurred more severely in the first group of mussels given daily. In other groups, liver damage decreased due to mussel consumption (Figure 1).




Figure 1. Control and experiment groups of liver tissue, H&E (CV: Central vein, long arrow: inflammation, short arrow: congestion, thick arrow: vacuolar degeneration).

NF- κB and TNF-α findings

Immunohistochemical staining results obtained from our study, we observed that NF-κB and TNF-α expression showed higher immunoreactivity in the liver due to the increase in the first experiment group, and staining was largely in the cell cytoplasm. The apoptotic mechanism was found to be very high in the liver tissues, especially in the first experimental group. Immunohistochemical staining with NF-κB, positive immunoreactivity was observed according to the amount of mussel given (Figure 2 and Figure 3). In the first experiment group, high reactivity was observed around the central vein. In the second and third groups, it was observed that the immunoreactivity around the central vein was moderate. A statistically significant difference was observed between the control and the first

experimental group of the subjects ($p<0.0001$). There was a weak significant difference between the control and third experiment group ($p<0.05$) (Table 2).




Figure 2. The immunohistochemical distribution of NF-κB in the control and experiment group of liver tissue (star and arrow: immunreactivity, CV: Central vein).





Figure 3. The immunohistochemical distribution of TNF-α in the control and experiment group of liver tissue (star and arrow: immunreactivity, CV: Central vein).

Table 2. TNF-α and NF-κB Immunoreactivty of control and all experiment groups. *p<0.0001 compared to the control group, **p<0.001 compared to the control group, ***p<0.05 compared to the control group. All experiment groups statistically significant differences were determined.



Heavy Metal Analysis Results

According to the research findings, considering the Cd, Pb, Cu, and Zn data in M. galloprovincialis, the places where the heavy metal concentration is the highest are Çamburnu, Yenikordon, and Çardak stations, respectively (Table 3).

Table 3. Heavy metal concentrations of *M. galloprovincialis* muscle tissue (µg/g dry weight.)

	Heavy metals			
Region	***Cd***	***Pb***	***Cu***	***Zn***
Çamburnu	1.22	0,56	1.35	18.74
Yenikordon	1.10	0,45	1.10	16.86
Çardak	0.84	0,42	0.72	16.24
Ortalama	1.05	0.47	1.05	17.28

There were differences and similarities between the groups in terms of histopathological damage in the liver tissues of the rats, which were collected with mussels collected from the Dardanelles. In our study, we researched Mediterranean mussels from shellfish, which have a high potential to cause liver damage to dietary habits. When mussels, which are both commercial and highly nutritious, are obtained from environmentally polluted environments, they can cause permanent damage to many organs, especially the digestive system (Gorinstein et al., 2008). Mussel dishes, which have become food culture especially in coastal areas and consumed frequently, can also cause permanent damage in one of the most important organs of our body, such as the liver. It supports this hypothesis in our research findings. The size of histopathological effects of mussels, which are frequently consumed and collected from unhealthy environments (exposed to industrial wastes), on the liver was found to be considerably large. Studies have reported that toxic substances such as heavy metals accumulate in fish and mussels due to marine pollution (Kayhan et al. 2006). In the researches, the presence of many heavy metals in single and bivalve seafood grown in the Dardanelles was determined (Ustunada et al., 2011). In our country, the annual pollution rate is quite high compared to the regions due to the transit ship passes of the Dardanelles and the Bosphorus. Again in the heavy metal detection studies carried out in the Mediterranean, trace elements such as cadmium, iron, and copper were found to accumulate in the bivalve (Göksu et al., 2005). Our findings show that, in the light of the data obtained from these studies, mussels can accumulate heavy metal and other pollutants in their tissues and create cytotoxic effects in humans and other organisms fed with them.

Long-term to heavy metals such as mercury (Hg), lead (Pb), chromium (Cr), cadmium (Cd), arsenic (As), copper (Cu), vanadium (V), nickel (Ni), and zinc (Zn) It has been reported that exposure may cause certain cancers in humans with chronic inflammation, cardiac, pulmonary and neurological effects (Mantovani et al., 2008; Nieboer et al., 2013). In our study, the detection of prominent foci of inflammation in the portal areas of the liver and parenchymal tissue was found similar to the findings of inflammation caused by the toxic effects of these investigators in people exposed to heavy metal. However, in our study, no findings related to liver cancer were encountered. Literature information has shown that heavy metals can cause cancer (Sá et al., 2016), but our study suggests that there is no cancer in the liver and that mussels are fed to rats for a short time, such as 30 days. Rats fed with mussels for a longer period may think that cancer may occur in the liver tissue.

Toxic liver diseases caused by drugs and chemicals mainly affect hepatocytes, bile ducts, vascular system, sinusoidal cells, and Kupffer cells, causing various morphological changes in the liver. Changing rates of hepatocyte damage, necrosis, lobular and portal inflammation are observed in hepatocellular damage. In its mildest form, balloon degeneration and apoptotic bodies, as well as focal necrosis foci, and inflammatory cells, including neutrophils and eosinophils, are observed in hepatocytes (Boone et al., 2005; David & Hamilton, 2010; Mantovani et al., 2008). In our study, balloon degeneration, portal and lobular fissures in hepatocytes are similar to the literature information seen in toxic liver cases. We detected necrosis, inflammation in the lobule and portal area. The fact that balloon degeneration is much more common in the group that feeds on mussels every day suggests that continuous mussel consumption causes more hepatocellular damage. In toxic hepatitis, cytotoxic hepatocellular damage can be seen as well as the cholestatic type (Bioulac-Sage & Balabaud, 2009; Lucena et al., 2008). Acute hepatocellular damage with cytotoxic effect may also be in mixed

form, where both cholestatic and cytotoxic damage can coexist. Cholestatic changes are accompanied by mild balloon degeneration, necrosis, and apoptosis (Schattenberg & Schuchmann, 2009). Our balloon degeneration and necrosis findings in our study suggest that mussels show cytotoxic effects in mixed form. We think that the source of histopathological damage occurring in the liver is due to the transition of heavy metals to mussels and toxic effects on the liver.

It has been shown that in liver injuries triggered by hemorrhagic shock, the protein supplements obtained from the crustaceans decrease the damage, and the level of inflammatory marker TNF α (Lee et al., 2012). However, it was shown that the severity of inflammation increased due to liver damage caused by shellfish collected and consumed without considering environmental pollution. Studies for its protective effects have also been shown in primary liver culture. When the protein products obtained from the crustaceans are given to the cells that are induced with carbon tetrachloride, it has been shown that the antioxidant level increases and the damage in the cells decreases (Chi et al., 2010). In another study, it was reported that pacific oyster extract decreased liver fibrosis, and inflammatory markers such as TGF-beta and NF-KB decreased (Zhou et al., 2015). In our study, it was determined that mussels caused degeneration of hepatocellular structure in the liver and that TNF-α immunoreactivity increased as the dose increased. TNF-α, a cytokine secreted for conservation purposes, shows that the liver's damaged structure can be cleaned by apoptosis and inflammatory events and re-trigger regeneration. When oxidative stress in tissues and associated damage to cells increases, activation of the TNF-alpha receptor is followed by activation of the nuclear factor kappa B (NF-kB). Thus, NF-κB goes to the nucleus, and then NF-κB activates genes that try to block TNF-induced apoptosis. In resting cells, NF-B is an inactive form in the cytoplasm (Karin, 2006; Schattenberg & Schuchmann, 2009). Our findings show that in previous studies, we observed that the expression of TNF-α and NF-κB increased in hepatocyte cytoplasms parallel to its regulatory role in liver tissue. As the hepatotoxic effect increased as consumption scallop increased, TNF-α and NF-κB expression was observed to be at the highest level in the first experimental group.

CONCLUSION

Our study results have confirmed the parameter results, which are encountered in many regions and countries, depending on the amount of consumption and environmental pollution. Heavy metal accumulation in mussels collected from the Dardanelles was observed to cause inflammation and degeneration in the liver. While searching for an alternative food source, environmental factors should not be ignored and it should be paid attention to the consumption of clean and healthy products since it causes tissue damage in many systems especially in the digestive system. Besides, before the consumption of mussels, especially heavy metal and other environmental pollution analysis should be done. Our people should be made aware of how much heavy metals they take with food accumulate in their organs and should be protected from possible liver diseases.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.

REFERENCES

Bioulac-Sage, P., & Balabaud, C. (2009). CHAPTER 40 - Toxic and Drug-Induced Disorders of the Liver. In R. D. Odze & J. R. Goldblum (Eds.), Surgical Pathology of the GI Tract, Liver, Biliary Tract, and Pancreas (Second Edition) (pp. 1059-1086). W.B. Saunders. https://doi.org/https://doi.org/10.1016/B978-141604059-0.50043-6

Boone, L., Meyer, D., Cusick, P., Ennulat, D., Bolliger, A., Everds, N., Meador, V., Elliott, G., Honor, D., Bounous, D., & Jordan, H. (2005). Selection and interpretation of clinical pathology indicators of hepatic injury in preclinical studies. Veterinary clinical pathology / American Society for Veterinary Clinical Pathology, 34, 182-188. https://doi.org/10.1111/j.1939-165X.2005.tb00041.x

Chang, L. W., Clarkson, T. W., Magos, L., & Suzuki, T. (1996). Toxicology of metals. Boca Raton (Fla.) : Lewis. http://lib.ugent.be/catalog/rug01:000695538

Chi, H. M., Chou, S. T., Lin, S. C., Su, Z. Y., & Sheen, L. Y. (2010). Protective effects of water extract of clam on normal and CCl_4-induced damage in primary cultured rat hepatocytes. Am J Chin Med, 38(6), 1193-1205. https://doi.org/10.1142/s0192415x10008561

David, S., & Hamilton, J. P. (2010). Drug-induced Liver Injury. US gastroenterology & hepatology review, 6, 73-80. https://pubmed.ncbi.nlm.nih.gov/21874146

Fergusson, J. E. (1990). The heavy elements: chemistry, environmental impact and health effects. Pergamon Press.

Figueras, A., Moreira, R., Sendra, M., & Novoa, B. (2019). Genomics and immunity of the Mediterranean mussel Mytilus galloprovincialis in a changing environment. Fish Shellfish Immunol, 90, 440-445. https://doi.org/10.1016/j.fsi.2019.04.064

Gezen, M.R. (2018). Hıstopathologıcal Changes In The Stomach Tıssue Of Rats Fed Wıth Mussels (Mytilus Galloprovincialis) Journal of Scientific Perspectives, 2, 2, 1-12. https://doi.org/10.26900/jsp.2018.14

Gorinstein, S., Leontowicz, M., Leontowicz, H., Namiesnik, J., Jastrzebski, Z., Drzewiecki, J., Park, Y.-S., Ham, K.-S., Heo, B.-G., & Trakhtenberg, S. (2008). Influence of mussels (Mytilus galloprovincialis) from polluted and non-polluted areas on some atherosclerosis indices in rats fed cholesterol. Food Chemistry, 111(2), 381-386. https://doi.org/https://doi.org/10.1016/j.foodchem.2008.03.082

Göksu, M. Z. L., Akar, M., Cevik, F., & Fındık, Ö. (2005). Bioaccumulation of some heavy metals (Cd, Fe, Zn, Cu) in two bivalvia species (Pinctada radiata Leach, 1814 and Brachidontes pharaonis Fischer, 1870). Turkish Journal of Veterinary and Animal Sciences, 29, 89-93.

Gürpınar, T., Ekerbiçer, N., Uysal, N., Barut, T., Tarakçı, F., & Tuglu, M. I. (2012). The effects of the melatonin treatment on the oxidative stress and apoptosis in diabetic eye and brain. ScientificWorldJournal, 2012, 498489. https://doi.org/10.1100/2012/498489

Hubbard, A. (2005). Heike B. Bradl, Editor, Heavy Metals in the Environment: Origin, Interaction and Remediation, Elsevier/Academic Press, London (2005) 269 pp. Journal of Colloid and Interface Science - J COLLOID INTERFACE SCI, 291, 307-307. https://doi.org/10.1016/j.jcis.2005.05.069

Karin, M. (2006). Nuclear factor-κB in cancer development and progression. Nature, 441(7092), 431-436. https://doi.org/10.1038/nature04870

Lee, R. P., Subeq, Y. M., Lee, C. J., Hsu, B. G., & Peng, T. C. (2012). Freshwater clam extract decreased hemorrhagic shock-induced liver injury by attenuating TNF-α production. Biol Res Nurs, 14(3), 286-293. https://doi.org/10.1177/1099800411408881

Lucena, M. I., García-Cortés, M., Cueto, R., Lopez-Duran, J., & Andrade, R. J. (2008). Assessment of drug-induced liver injury in clinical practice. Fundam Clin Pharmacol, 22(2), 141-158. https://doi.org/10.1111/j.1472-8206.2008.00566.x

Mantovani, A., Allavena, P., Sica, A., & Balkwill, F. (2008). Cancer-related inflammation. Nature, 454(7203), 436-444. https://doi.org/10.1038/nature07205

Nieboer, E., Tsuji, L. J., Martin, I. D., & Liberda, E. N. (2013). Human biomonitoring issues related to lead exposure. Environ Sci Process Impacts, 15(10), 1824-1829. https://doi.org/10.1039/c3em00270e

Öztürk, Ş., Kılıçaslan Sönmez, P., Özdemir, İ., Topdağı, Y., & Tuğlu, M. (2019). Kemik iliği kaynaklı mezenkimal kök hücrelerin deneysel Asherman modeli üzerine antiapoptotik ve proliferatif etkisi. Cukurova Medical Journal, 44, 434-443. https://doi.org/10.17826/cumj.573200

Palmes, D., & Spiegel, H.-U. (2004). Animal models of liver regeneration. Biomaterials, 25(9), 1601-1611. https://doi.org/https://doi.org/10.1016/S0142-9612(03)00508-8

Pourret, O., & Hursthouse, A. (2019). It's Time to Replace the Term "Heavy Metals" with "Potentially Toxic Elements" When Reporting Environmental Research. International journal of environmental research and public health, 16(22), 4446. https://doi.org/10.3390/ijerph16224446

Sá, I., Semedo, M., & Cunha, M. (2016). Kidney cancer. Heavy metals as a risk factor. Porto Biomedical Journal, 1, 25-28. https://doi.org/10.1016/j.pbj.2016.03.006

Schattenberg, J. M., & Schuchmann, M. (2009). Diabetes and apoptosis: liver. Apoptosis, 14(12), 1459-1471. https://doi.org/10.1007/s10495-009-0366-2

Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., & Sutton, D. J. (2012). Heavy metal toxicity and the environment. Experientia supplementum (2012), 101, 133-164. https://doi.org/10.1007/978-3-7643-8340-4_6

Ustunada, M., Erduğan, H., Yılmaz, S., Akgul, R., & Aysel, V. (2011). Seasonal concentrations of some heavy metals (Cd, Pb, Zn, and Cu) in Ulva rigida J. Agardh (Chlorophyta) from Dardanelles (Canakkale, Turkey). Environmental Monitoring and Assessment, 177(1), 337-342. https://doi.org/10.1007/s10661-010-1637-7

Voudanta, E., Kormas, K. A., Monchy, S., Delegrange, A., Vincent, D., Genitsaris, S., & Christaki, U. (2016). Mussel biofiltration effects on attached bacteria and unicellular eukaryotes in fish-rearing seawater. PeerJ, 4, e1829-e1829. https://doi.org/10.7717/peerj.1829

Wang, S., & Shi, X. (2001). Molecular mechanisms of metal toxicity and carcinogenesis. Mol Cell Biochem, 222(1-2), 3-9.

Yang, Y. M., & Seki, E. (2015). TNFα in liver fibrosis. Current pathobiology reports, 3(4), 253-261. https://doi.org/10.1007/s40139-015-0093-z

Yigit, M., Celikkol, B., Yilmaz, S., Bulut, M., Ozalp, B., Dwyer, R. L., Maita, M., Kizilkaya, B., Yigit, Ü., Ergün, S., Gürses, K., & Buyukates, Y. (2018). Bioaccumulation of trace metals in Mediterranean mussels (Mytilus galloprovincialis) from a fish farm with copper-alloy mesh pens and potential risk assessment. Human and Ecological Risk Assessment: An International Journal, 24(2), 465-481. https://doi.org/10.1080/10807039.2017.1387476

Zhou, J., Liang, Y., Pan, J. X., Wang, F. F., Lin, X. M., Ma, R. J., Qu, F., & Fang, J. Q. (2015). Protein extracts of Crassostrea gigas alleviate CCl_4-induced hepatic fibrosis in rats by reducing the expression of CTGF, TGF-β1 and NF-κB in liver tissues. Mol Med Rep, 11(4), 2913-2920. https://doi.org/10.3892/mmr.2014.3019

Acta Aquatica Turcica

Makale Sayısı	Atıf Sayısı	Kendine Atıf Sayısı
321	339	37

Atıf Alan Makale Sayısı	Atıf Ortalaması	Kendine Atıf Oranı
141	1,06	%10,91


Konu Kategorisi: Fen > Temel Bilimler


Konu Alanları: Zooloji, Balıkçılık



EISSN:	2651-5474
İlk Yayın Yılı:	1988
Dizinlendiği Yıllar:	2018-2025 (Fen)
Yayın Periyodu:	Mart, Haziran, Eylül, Aralık
Yayın Dili:	Türkçe İngilizce
Editör:	Yunus Ömer BOYACI
Yayıncı:	Yunus Ömer BOYACI
Web Adresi:	http://dergipark.org.tr/actaquatr


Makale & Atıf Sayısı






"Grafik Verisi" ©2026 - TR Dizin

Makale Türleri





"Grafik Verisi"©2026 - TR Dizin

Derginin Makaleleri | **Derginin Aldığı Atıflar**

Türk Marketlerinde Satışa Sunulan Çeşitli Balık Ürünlerinin Biyojenik Amin ve Trimetilamin İçerikleri
Ali Serhat ÖZKÜTÜK, Gülsün ÖZYURT, Esmeray Kuley
MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 13 - 23
Analiz Listesine Ekle
Öz

Silurus triostegus'un Gonat Dokusu Yağ Asidi İçeriğinin Mevsimsel Değişimi
Semra KAÇAR, Mehmet BAŞHAN
MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 24 - 37
Analiz Listesine Ekle
Öz
1 Atıf

Assessment of Sediment Dredging Effectiveness by Considering Overlying Water and Pore Water Heavy Metals in Lake Mogan, Turkey
Arzu BİNİCİ, Serap PULATSÜ
MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 38 - 48
Analiz Listesine Ekle
Öz
1 Atıf

Farklı Zenginleştirici Emülsiyonların Rotifer (Brachionus plicatilis) Yığın Kültüründe Kullanımı
HASAN BATUHAN EMRE OZDOGAN, Sevgi SAVAŞ
MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 49 - 59
Analiz Listesine Ekle
Öz

Kuzey Ege Denizi'nde Uzatma Ağlarında Farklı Ağ İpi Kalınlığının Av Verimi ve Av Kompozisyonu Üzerine Etkisi
Mustafa BAKIRCI, Adnan AYAZ, Alkan ÖZTEKİN, Gençtan Erman UĞUR
MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 60 - 75
Analiz Listesine Ekle
Öz

Ammothella longioculata (Arthropoda, Pycnogonida): A Sea Spider New For the Mediterranean Coast of Turkey
Cengiz KOÇAK
2

Atıf

MAKALE
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 76 - 80
Analiz Listesine Ekle
Öz

MAKALE
Effects of Thyme and Thyme Oil on Growth of White Shrimp, Litopenaeus vannamei
Mevlüt AKTAŞ, Münevver Ayçe GENÇ, Yasemin BİRCAN YILDIRIM, Doğukan KAYA, Özlem ÇALIŞICI NARİN, Ercüment GENÇ
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 81 - 92
Analiz Listesine Ekle
Öz

MAKALE
Çanakkale'nin Yenice ve Bayramiç İlçelerinde Su Ürünleri Tüketiminin Değerlendirilmesi
Kahraman SELVİ, Burcu KAYA, Seda ÖZDİKMENLİ TEPELİ, Gülcan KANDEMİR
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 93 - 108
Analiz Listesine Ekle
Öz
7
Atıf

MAKALE
Heavy Metal Contamination Status and Risk Assessment in Surface Sediments of Köyceğiz Lagoon Estuary System (KLES) (South-West Anatolia)
Murat YABANLI, AYKUT YOZUKMAZ, İdris ŞENER, Tülin ÇOKER, Hatice H. YAPICI, Esra ÇETİN
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 109 - 120
Analiz Listesine Ekle
Öz
2
Atıf

MAKALE
Extended North Distribution of Penicillus capitatus Lamarck 1813 off the Turkish Shores
Levent ÇAVAŞ
Acta Aquatica Turcica
Yıl: 2022 Cilt: 18 Sayı: 1 Sayfa: 121 - 130
Analiz Listesine Ekle
Öz

Toplam Kayıt: 321

1 / 33 10

İLETİŞİM
TÜBİTAK ULAKBİM TR Dizin
Yüzüncüyıl, İşçi Blokları Mahallesi
Muhsin Yazıcıoğlu Caddesi No:51/C
06530 Çankaya / ANKARA +90 (312) 298 92 00
trdizin@tubitak.gov.tr

T.C. SANAYİ VE TEKNOLOJİ BAKANLIĞI
#MİLLİ TEKNOLOJİ HAMLESİ
TÜBİTAK ULAKBİM