**FEB - FRESENIUS ENVIRONMENTAL BULLETIN**
Founded jointly by F. Korte and F. Coulston
Production by PSP - Vimy Str. 1e, 85354 Freising, Germany in
cooperation with PRT-Parlar Research & Technology
Vimy Str 1e, 85354 Freising
Copyright© by PSP and PRT, Vimy Str. 1e, 85354 Freising, Germany
All rights are reserved, especially the right to translate into foreign language or other processes - or convert to a machine language, especially for data processing equipment - without written permission of the publisher. The rights of reproduction by lecture, radio and television transmission, magnetic sound recording or similar means are also reserved.
**Printed in Germany-ISSN 1018-4619**

**FEB-EDITORIAL BOARD**

**CHIEF EDITOR:**
**Prof. Dr. H. Parlar**
Parlar Research & Technology-PRT
Vimy Str.1e
85354 Freising, Germany

**MANAGING EDITOR:**
**Dr. P. Parlar**
Parlar Research & Technology
PRT, Vimy Str.1e
85354 Freising, Germany

**CO-EDITORS:**

***Environmental Spectroscopy***
**Prof. Dr. A. Piccolo**
Universita di Napoli "Frederico II"
Dipto. Di Scienze Chemica Agrarie
Via Universita 100, 80055 Portici, Italy

***Environmental Biology***
**Prof. Dr. G. Schuurmann**
UFZ-Umweltzentrum
Sektion Chemische Ökotoxikologie
Leipzig-Halle GmbH,
Permoserstr.15, 04318
04318 Leipzig, Germany
**Prof. Dr. I. Holoubek**
Recetox-Tocoen
Kamenice126/3, 62500 Brno, Czech Republic
**Prof. Dr. M. Hakki Alma**
Igdir Universitesi
76000, Igdir, Turkey
**Prof. Dr. A. Reichlmayr-Lais**
Technical University of Munich
Arcisstraße 31
80333 Muenchen,Germany

***Environmental Management***
**Dr. K. I. Nikolaou**
Env.Protection of Thessaloniki
OMPEPT-54636 Thessaloniki
Greece

***Environmental Toxicology***
**Prof. Dr. H. Greim**
Senatkommision – DFG / TUM
85350 Freising, Germany

***Environmental Proteomic***
**Dr. A. Fanous**
Halal Control GmbH
Stahlstraße 44
D-65428 Rüsselsheim, Germany

***Environmental Analytical Chemistry***
**Prof. Dr. M. Bahadir**
Lehrstuhl für Ökologische Chemie
und Umweltanalytik
TU Braunschweig
Lehrstuhl für Ökologische Chemie
Hagenring 30, 38106 Braunschweig, Germany
**Dr. D. Kotzias**
Via Germania29
21027 Barza(Va), Italy
**Prof. Dr. R. Kallenborn**
Norwegian University of Life Sciences
Universitetstunet 3
1430 As,Norway

***Environmental Education***
**Prof. Dr. C. Bayat**
Yeni Yüzyil Üniversitesi
34010 Zeytinburnu, Istanbul, Turkey

***Environmental Medicine***
**Prof. Dr. I. Tumen**
Bandirma 17 Eylül Üniversitesi
10200 Bandirma, Turkey
**Dr. J. Burhenne**
Universitaet Klinikum
Im Neuenheim Feld 410
69120 Heidelberg,Germany

*Advisory Board*
**K. Bester, K. Fischer, DCG. Muir,**
**R. Niessner,W.Vetter, D. Steinberg,**
**J. P. Lay, L. O. Ruzo**

*Marketing Manager*
**Cansu Ekici, MSC. of B.A.**
PRT-Research and Technology
Vimy Str 1e
85354 Freising, Germany
**E-Mail: parlar@wzw.tum.de**
**parlar@prt-parlar.de**
**Phone: +49/8161887988**

**Fresenius Environmental Bulletin is abstracted/indexed in:**
**Biology & Environmental Sciences, BIOSIS, CAB International, Cambridge Scientific abstracts, Chemical Abstracts, Current Awareness, Current Contents/Agriculture, CSA Civil Engineering Abstracts, CSA Mechanical & Transportation Engineering, IBIDS database, Information Ventures, NISC, Research Alert, Science Citation Index (SCI), Scisearch, Selected Water Resources Abstracts**

# CONTENTS

## ORIGINAL PAPERS

HYPERSPECTRAL PREDICTION OF SOIL ORGANIC MATTER IN WINTER WHEAT FIELD BASED ON CONTINUUM-REMOVED METHOD 7979
**Chao Wang, Xingxing Qiao, Zhuoya Liang, Guangxin Li, Lujie Xiao, Meijun Zhang, Xiaoyan Song, Muhammad Saleem Kubar, Meichen Feng, Wude Yang***

PHYSIOLOGICAL EFFECT OF THE COMBINED STRESS OF FREEZE-THAW AND SALINE-ALKALI ON LEYMUS CHINENSIS RHIZOMES 7987
**Guozhang Bao, Yan Qu, Bairu Yan, Xiaoting Liu, Mengyu Zhang, Wenyi Tang, Wanyu Dong**

A NEW LOGGING INTERPRETATION METHOD FOR DESCRIBING THE COMPRESSIVE AND SHEAR STRENGTH OF UNDERGROUND TIGHT SANDSTONE 7996
**Ping Liu, Lu Wang, Shuai Yin**

CORRELATION AND PATH COEFFICIENT ANALYSIS FOR COTTON (*GOSSYPIUM HIRSUTUM* L. ) IN DIFFERENT SOIL TEMPERATURES 8004
**Hatice Cokkizgin**

EVALUATION OF ECOLOGICAL FUNCTIONS OF URBANPARK LANDSCAPES BASED ON ANALYTIC HIERARCHY PROCESS METHOD 8013
**Youwen Wang, Yan Zhang**

DETERMINATION OF NUTRITIONAL EVALUATION FATTY ACID COMPOSITION AND GROWTH KINETICS OF *SPIROGYRA* STRAINS 8021
**Ilkay Acikgoz Erkaya, Dilek Yalcin**

INFLUENCING FACTORS AND OPTIMIZATION WAYS OF INTERNAL CONTROL ENVIRONMENT IN ANIMAL HUSBANDRY ENTERPRISES 8034
**Lu Tang, Shufei Wang, HongYan Deng, ZongQian Li, XuHai Wang**

STUDY ON ENTERPRISES' GREEN DEVELOPMENT UNDER THE BP ARTIFICIAL NEURAL NETWORK METHODS BASED ON ECOLOGICAL PERSPECTIVE 8041
**Lu Zhang, Hao Zhang, Danting Zheng, Ronghua Chang, Sanggyun Na**

STUDIES ON SURVEILLANCE AND PATHOLOGY OF *E. COLI* INFECTIONS IN POULTRY IN SINDH PROVINCE OF PAKISTAN 8051
**Zubair Ahmed Soomro, Zaheer Ahmed Nizamani, Nazeer Hussain Kalhoro, Shakeel Ahmed Soomro, Haiming Yu, Kunjie Chen, Mansoor Tariq, Waseem Ali Vistro, Hafeez-ur-Rehman Mangio, Anees Ahmed Soomro**

ANALYSIS ON MAIN FACTORS AFFECTING THE QUALITY OF MICROBIOLOGICAL EXAMINATION AND DRUG RESISTANCE OF PATHOGENS 8063
**Weiping Du, Rui Zhang, Zhixi Hu, Sirong Mi, Ningmei Zhang, Lifen Bai, Yongxia Liu**

ELIMINATION OF SUSPENDED MATERIALS IN IRRIGATION WATER WITH DIFFERENT MEDIA FILTERS 8070
**Ozlem Korkut, Sultan Kiymaz, Yasemin Kuslu**

MECHANISM ANALYSIS OF MODIFIED RICE HUSK GASIFICATION CHAR TO REMOVE MERCURY BY ADSORPTION KINETICS 8078
**Ru Yang, Yongfa Diao, Yaolin Lou**

ANTIMICROBIAL ACTIVITY OF DIFFERENT SPICE HERBS EXTRACTS AGAINST SOME PATHOGENIC AND NON PATHOGENIC STRAINS OF E. COLI: IN VITRO - AND IN VIVO - STUDY 8084
**Sadia Alam, Aisha Bibi, Muhammad Mumtaz Khan, Nouman Rashid Siddique, Samina Khalil, Najma Ayub**

ANALYSIS OF HERPETOFAUNAL DIVERSITY OF ISTANBUL ISLANDS WITH THE NEW RECORDS OF THREE LOCALITIES 8095
**Nilgun Kaya, Murat Tosunoglu**

RESEARCH ON THE TEACHING BEHAVIOR OF PHYSICAL EDUCATION IN THE VIEW OF ECOLOGICAL CLASSROOM 8100
**Chao Liu, Cuixiang Dong**

STUDY ON SATURATED HYDROCARBONS IN ORGANIC RICH ROCKS INDICATIVE SIGNIFICANCE OF SPECIAL COMPOUND MOLECULES FOR HYDROCARBONS 8109
**Shaochun Wang, Zengyuan Zhou, Dexiang Yang, Shuguang Chen, Jianping Wu, Wenjun Zhang, Jing Liu, Huilai Wang**

BIOREMEDIATION OF ARSENIC IN RICE BY ARSENIC RESISTANT BACTERIA ISOLATED FROM LAKE VAN, TURKEY, THE LARGEST SODA LAKE 8117
**Asli Bayer, Esra Ersoy Omeroglu**

CHARACTERISTICS OF AIR POLLUTION IN JILIN CITY AND ITS EFFECTS ON THE RESPIRATORY SYSTEM OF OUTDOOR EXERCISERS 8130
**Songbo He, Hongchun Jia**

ISOLATION AND MORPHO-MOLECULAR IDENTIFICATION OF INDIGENOUS SOIL-BORNE ENTOMOPATHOGENIC FUNGI, AND THEIR PATHOGENICITY TO GALLERIA MELLONELLA L. (LEPIDOPTERA: PYRALIDAE) AS A MODEL INSECT 8138
**Derya Baki, Fedai Erler, Hilal Sule Tosun, Baris Imrek**

INDOOR CUCUMBER PEST MONITORING BASED ON CHLOROPHYLL FLUORESCENCE SPECTRUM ANALYSIS 8149
**Wenyi Chen, Xinghui Zhu**

PHOTOSENSITIZING EFFECTS OF HEMATOPORPHYRIN DIHYDROCHLORIDE AGAINST THE RED PALM WEEVIL RHYNCHOPHORUS FERRUGINEUS (COLEOPTERA: CURCULIONIDAE) 8158
**Zeinab Abdel-Moaty, Salah Abdelsalam**

STUDY ON THE DIAGENESIS LOWER LIMIT OF THE RELATIVELY HIGH-QUALITY PALAEOGENE RESERVOIRS IN THE XIHU DEPRESSION 8165
**Linhai Zhao, Qing Miao, Guosheng Xu, Yong Liu, Changzhi Li, Zhixing Zhao**

ANALYSIS OF THE IMPACT OF DRILLING MUD SOLIDIFICATION TREATMENT ON SOIL ENVIRONMENT 8179
**Ming Yi, Hao Huang, Bingqing Li, Bing Liu, Zhiqiang Huang**

FLOCCULATION AND REMOVAL OF MICROCYSTIS AERUGINOSA BY THE NATURAL COAGULANT FROM CITRUS PEEL 8187
**Linlin Cai, Hongqiang Wang, Shengyuan Yuan**

EFFECTS OF PROPYLENE GLYCOL ON OVARIAN RESTORATION, REPRODUCTIVE PERFORMANCE, METABOLIC STATUS AND MILK PRODUCTION DURING TRASITION PERIOD OF EWES IN SUBTROPICS 8192
**Abd El-Nasser Ahmed Mohammed, Fahad Al-Hizab, Shaker Al-Suwaiegh, Tarek Alshaheen, Ayman Kassab, Hatem Hamdon, Waleed Senosy**

EFFECTS OF FEEDING GREEN MICROALGAE AND *NIGELLA SATIVA* ON PRODUCTIVE PERFORMANCE AND METABOLIC PROFILE OF BOER GOATS DURING PERIPARTUM PERIOD IN SUBTROPICS 8203
**Ali M Ali, Tarek Alshaheen, Waleed Senosy, Abd El-Nasser Mohammed, Ayman Kassab**

RESEARCH ON WATER DEMAND PLANNING OF URBAN LANDSCAPE ECOLOGICAL ENVIRONMENT 8213
**Qiwei Liu**

TiO2/Fe-ZSM5 PHOTOCATALYTIC DEGRADATION OF PRINTING AND DYEING WASTEWATER 8221
**Endong Wang**

THE CARBON AND OXYGEN ISOTOPE CHARACTERISTICS AND PALEOENVIRONMENTAL SIGNIFICANCE OF LOWER–MIDDLE CARBONIFEROUS CARBONATE ROCKS OF THE EASTERN CASPIAN BASIN 8228
**Bobiao Liu, Lun Zhao, Chengqian Tan, Huiyuan Bian, Yinchao Huai, Yaping Lin, Xiangliang Qiu**

FOREST SITES MAPPING IN SERBIA 8244
**Olivera Kosanin, Marko Perovic, Milan Knezevic, Rade Cvjeticanin, Janko Ljubicic**

MICROBIOLOGICAL MONITORING OF WATER SUPPLY BY INTESTINAL ENTEROCOCCI USING THE MEMBRANE FILTRATION METHOD 8252
**Safta Victor Viorel, Ciuca Viviana Carmen, Voicu Gheorghe**

DEFICIT IRRIGATION AS SUSTAINABLE PRACTICE WITH VARYING SOWING WINDOWS FOR ENHANCING WATER USE EFFICIENCY, GRAIN QUALITY AND BIOCHEMICAL CHARACTERISTICS OF CHINESE WHEAT HYBRID LINES IN COMPARISON TO LOCAL CHECK UNDER SEMI- ARID CLIMATIC CONDITION 8260
**Muhammad Mehran Anjum, Muhammad Arif**

LAND USE COVER CHANGE ANALYSIS (1985-2020) OF AYVALIK DISTRICT BALIKESIR TURKEY USING GEOGRAPHIC INFORMATION SYSTEMS AND REMOTE SENSING 8278
**Figen Altiner, Abdullah Kelkit**

EVALUATION OF SURFACE WATER QUALITY IN PANJIN OF LIAO RIVER BASIN BY PRINCIPAL COMPONENT ANALYSIS 8284
**Lei Chao, Yu Cao, Su Chen, Yao Wang, Yafeng Li**

GENETIC CHARACTERIZATION OF *PSEUDOMONAS SAVASTANOI* PV. *SAVASTANOI* STRAINS ISOLATED FROM DIFFERENT OLIVE CULTIVARS GROWN IN JORDAN BY PCR-RFLP AND AFLP 8292
**Bilal Ibrahim Wreikat**

STUDY ON THE INFLUENCE OF ROCK STRENGTH ANOMALIES ON THE COLLAPSE PRESSURE OF THE BOREHOLE WALL AND ITS INSTABILITY AREA DURING OIL AND GAS DEVELOPMENT 8306
**Hui Tian**

ASSESSMENT OF PHYSICOCHEMICALS AND MICROBIAL PARAMETERS OF DIFFERENT SUPPLIES OF WATER IN KOH-E-SULEMAN DERA GHAZI KHAN PUNJAB, PAKISTAN 8315
Shan Ali Raza[1], Naeem Hussain[2], Asad Ullah[3], Mohammed Shah Jamal[4], Kuaanan Techato[1,5,*]

DETERMINATION OF HEAVY METALS IN HONEY BY ATOMIC ABSORPTION SPECTROMETRY 8325
**Albana R Mehmeti, Musaj Pacarizi, Ismet Hashani, Majlinda Daci-Ajvazi, Sevdije Govori, Albert Maxhuni**

EFFECT OF HUMIC ACID ON THE INFILTRATION RESISTANCE OF SOIL WATER 8329
**He Jing, Dong Qiguang**

THE CONSTRUCTION OF REAL-TIME DETERMINATION SYSTEM AND UNCERTAINTY EVALUATION FOR THE EFFICENCY OF VOLATILE ORGANIC COMPOUNDS PHOTOCATALYSIS DECOMPOSITION 8336
**XingJie Lu, Ke Zhang, YongHong Zhu, WenSi Liu, JiaCheng Fan, XiaoWei Wang**

KEY DRILLING TECHNOLOGY OF 4000M ULTRA LONG HORIZONTAL WELL IN TIGHT GAS RESERVOIR OF CHANGQING OILFIELD, CHINA 8343
**Huajun Meng, Shuhui Wang, Chenhui Yang, Liangjun Wu, Yongjun Chen, Fang Cai**

MORPHOLOGICAL CHARACTERISTICS OF CONTAINERIZED AND BARE-ROOT SEEDLINGS IN *JUNIPERUS EXCELSA* AND *J. FOETIDISSIMA* 8351
**Yunus Eser**

ANALYSIS ON THE CHARACTERISTICS OF ACID RAIN AND THE CORRELATION OF MAIN ACID GAS INFLUENCING FACTORS IN HENGYANG CHINA 8356
**Zuoyun Liu, Henglin Liu**

METAL LEVELS OF MACROALGAE FROM ISKENDERUN AND MERSIN BAY TURKEY EASTERN MEDITERRANEAN 8365
**Sibel Alagoz Erguden, Deniz Erguden, Nuray Ciftci, Hasan Deniz Akbora, Deniz Ayas**

EVALUATION ON THE COORDINATED DEVELOPMENT OF AGRICULTURAL ECONOMY AND RURAL ECOLOGICAL ENVIRONMENT IN ANHUI PROVINCE 8372
**Minmin Deng**

COMPREHENSIVE EVALUATION OF ECOLOGICAL SECURITY OF ANHUI PROVINCE 8382
**Gang He, Jun Ruan, Keyu Bao, Yinging Wang, Yangqiu Zhao**

EXPERIMENTAL RESEARCH ON THE HTHP RHEOLOGICAL PROPERTIES OF ULTRA-HIGH TEMPERATURE AND ULTRA-HIGH PRESSURE WATER BASED DRILLING FLUID 8392
**Wei Gao, Yinting Li, Fuchun Yu, Junwei Fang, Shuanggui Li, Hui Mao**

ANALYSIS ON THE ECONOMIC ECOLOGICAL ENVIRONMENT AND SOCIAL EFFECTS OF RURAL TOURISM DEVELOPMENT 8404
**Naipeng Bu, Jianyang Guo, Rachel Lam, Haiyan Kong, Shun Yi, Yanan Li**

EXPERIMENTAL STUDY ON RHEOLOGICAL PROPERTIES OF ULTRA-HIGH TEMPERATURE WATER BASED DRILLING FLUID FOR GEOTHERMAL WELL DRILLING APPLICATIONS 8411
**Yinde Zhang, Hui Mao, Junwei Fang, Xianbin Zhang, Xinxin Wen**

IDENTIFICATION AND CHARACTERIZATION OF POLYETHYLENE-DEGRADING MICROORGANISMS IN THE INTESTINAL TRACT OF LARVAE OF THE GREATER WAX MOTH, *GALLERIA MELLONELLA* L. 8423
**Minmin Mao, Yafei Liu, Yibin Cao, Xia Liu, Xuexiang Weng, Yongming Ruan**

EFFECTS OF AMMONIUM - AND NITRATE - NITROGEN ON THE GROWTH OF BLUEBERRY CLUSTER BRANCHES 8432
**Yinan Ding, Lihui Zhang, Kai Song, Lijiao Zhang, Juan Yan, Haijiao Wang, Siqi Liu, Fei Lv, Guangye Wang**

DEBRIS FLOW RISK ASSESSMENT MODEL BASED ON LINEAR PROGRAMMING OPTIMIZATION ALGORITHM UNDER GEOLOGICAL SAFETY ENVIRONMENT 8439
**Shixin Zhang, Li Li, Yue Qiang, Dongsheng Zhao, Changsi Xia, Zhou Zheng**

RESERVOIR CONFIGURATION ANATOMY OF BRAIDED RIVER DELTA IN Es2 OF YING-14 FAULT BLOCK IN DONGXIN OILFIELD CHINA 8447
**Ruixiang Zhang, Liang Gao, Qinglin Bai, Xuejuan Zhu**

RESERVOIR CHARACTERISTICS AND HIGH-QUALITY RESERVOIR PREDICTION OF THE CARBONIFEROUS KARASAYI FORMATION IN TAHE OILFIELD 8458
**Hongtao Ma, Honggang Liang, Li Sun, Feng Deng, Wei Xian, Rui Guo**

EXPERIMENTAL STUDY ON STRESS SENSITIVITY OF PERMEABILITY AND ELASTIC PARAMETERS OF UNCONSOLIDATED SANDSTONE RESERVOIRS 8467
**Xueyou Zhang**

PERFORMANCE EVALUATION OF COMPOSITE WATER SOLUBLE TEMPORARY PLUGGING AGENT SYSTEM FOR OIL WELL FLUSHING 8476
**Hongwei Yu**

STUDY ON ECOLOGICAL DAMAGE ASSESSMENT AND RESTORATION OF COASTAL ZONE BASED ON THE PSR/RS MODE 8484
**Ruizhen Zhao**

A NOVEL DYNAMIC PLANNING METHOD FOR RESCUE ROUTES OF REFUGE BUILDINGS IN RESPONSE TO GEOLOGICAL ENVIRONMENTAL DISASTERS BASED ON SPATIAL LAYOUT THEORY 8491
**Yang Li, Ping Liu**

STUDY ON THERMAL INSULATION PERFORMANCE EVALUATION AND PROCESS PARAMETERS OPTIMIZATION OF GREEN BUILDING WALL 8498
**Xin Ye**

LAND USE CHANGE AND ECOLOGICAL ENVIRONMENT EFFECTS IN COAL RESOURCE CITY: A CASE STUDY OF PINGDINGSHAN, HENAN PROVINCE 8506
**Guangfang Xiang, Rong Hua**

EVALUATION OF URBAN LANDSCAPE ECOLOGICAL RISK BASED ON THE THEORY OF LANDSCAPE ECOLOGY 8514
**Chao Chen, Pangong Cao**

INVESTIGATING THE EFFECT OF *SPİRULİNA* ON THE HEMOXYGENASE RELEASE IN THE UTERUS TISSUE OF THE EXPERIMENTAL THYROTOXICOSIS- INDUCED RATS 8521
**Nilgun Atam, Sevda Elis Yildiz**

DETERMINATION OF SPRAY PERFORMANCE OF NEW AND WORN FLAT FAN NOZZLES AND SPRAY ANGLE BASED ON NOVEL IMAGE ANALYSIS SOFTWARE 8532
**Ali Bolat, Omer BarisOzluoymak, Ali Bayat**

RESEARCH REGARDING IMPACT OF INDOOR NOISE EXPOSURE ON YOUNG ADULTS 8541
**Aida Sahmurova, Nilgun Balkaya, Gul Alpar**

CARRYING CAPACITY AND THE EFFECT OF BATHERS ON THE SANITARY QUALITY OF WATER AT A RURAL RECREATIONAL BEACH 8549
**Claudia Esmeralda Leon-Lopez, Gisela Garcia-Morales, Jorge Eduardo Chavez-Villalba, Renato Arturo Mendoza-Salgado, Jaqueline Garcia-Hernandez, Lia Celina Mendez-Rodriguez, Jose Alfredo Arreola-Lizarraga**

APPLICATION OF TWO-STEP POD REDUCED ORDER MODEL IN NUMERICAL SIMULATION OF OIL RESERVOIR 8559
**Jing Cao**

NOVEL ADSORPTION MATERIALS BASED ON GRAPHENE OXIDE/DIATOMITE COMPOSITE MATERIALS AND THEIR ADSORPTION PERFORMANCE FOR METHYLENE BLUE 8566
**Yinghu Zhao, Li Gao, Linbian Wang, Shuli Zhang, Jun Xie**

MINIMIZING THE LOST CIRCULATION IN OIL-BASED DRILLING FLUIDS USING OIL ABSORBING EXPANSIVE RUBBER 8576
**Baocheng Wu, Shengjiang Xu, Cheng Ye, Kecheng Liu, Chao Ma, Danfeng Jia**

RANGE VEGETATION ANALYSIS OF PAKISTAN'S SOUTHERN PUNJAB'S RANGELAND RAKH CHOTI DALANA 8588
**Muhammad Yasin, Muhammad Zubair, Syed Bilal Hussain, Ahsan Ul Haq, Ahmad Hussain, Long Wenxing**

DISTRIBUTION CHARACTERISTICS OF PALEOPRESSURE IN THE LOWER PALEOZOIC OF THE ORDOS BASIN, CHINA 8595
**Xinshe Liu, Rizhou Cao, Xing Gao, Zixing Lu**

MODELLING AND ANALYSIS OF WATER QUALITY FOR RESERVOIR IN THE NORTHERN COLD REGION OF CHINA 8602
**Wei Su, Guozheng Wu, Gula Tang, Shihui Shi**

ENVIRONMENTAL HETEROGENEOUS DATA ANALYSIS OF WATER QUALITY TO TRACE POTENTIAL FACTORS AND IDENTIFY SOURCES IN AN URBANIZED RIVER 8613
**Kuotian Lu, Hongjie Gao, Huibin Yu, Dongping Liu, Fangyuan Jin, Ningmei Zhu, Siyu Wang**

STUDY ON THE CHEMICAL COMPOSITION ANALYSIS FROM GREEN PINE ALGAE 8624
**Na Li, Yunyun Gai, Ling Su, Yanli Chen**

QUANTIFICATION AND HEALTH RISK ASSESSMENT OF TWENTY-FIVE ELEMENTS IN CHEESE, BEEF, MUTTON AND LIVER PRODUCED IN NUBA MOUNTAINS, SUDAN 8630
**Abdelbagi N A Elrashid, Abubakr M Idris, Ibrahim M Ahmed, Ali H Ali, Mohammed A H Eltayeb**

PHYTOCHEMICAL SCREENING OF AMARANTH HERBAGE AND THEIR IMPRESSION ON THE ANTIOXIDANT ACTIVITY WITH AN ASSESSMENT OF FEEDING QUALITY 8641
**Nafiz Celiktas, Ersin Can, Ibrahim Atis, Derya Durak**

PRESERVATION OF VITAMIN-E AGAINST NEPHROTOXIC EFFECT INDUCED BY SUBACUTE DICHLORVOS APPLICATION 8651
**Gokhan Nur, H Ahmet Deveci, Evren Koc**

INFLUENCE OF SOME HERBICIDES ON WEEDS QUALITY AND PRODUCTIVITY OF POTATO CROP (*SOLANUM TUBEROSUM* L.) IN NEWLY RECLAIMED AREAS 8660
**Amin Abd El-Baky Zein, Mohammad Abd El-Salam Abd El-Baky, Aly Aly Hassan Sharshar, Abdulla basher Mohammad El-badawy**

EVALUATION AND DEVELOPMENT TREND PREDICTION OF GUANGDONG'S ELECTRIC ENERGY AND ENVIRONMENTAL EFFICIENCY 8668
**Junjun Zheng, Yongji Wang**

PROTECTION PRIORITY PLANNING OF TOURISM AND AGRICULTURE WITHIN THE SCOPE OF SUSTAINABLE DEVELOPMENT BARTIN KIZILKUM-MUGADA COASTAL REGION 8675
**Pinar Bollukcu**

AN INVESTIGATION ON GYRODACTYLID HELMINTH FAUNA OF CHUB, SQUALIUS CEPHALUS (LINNAEUS, 1758) IN RELATION TO SEASONAL CHANGES, HOST AGES AND SEX 8685
**Huseyin Yazmen, Mehmet Oguz Ozturk**

SEASONAL VARIATION OF LAMPARELLI TROPHIC STATE INDEX (TSI) IN KUNE -VAINI LAGOON SYSTEM 8690
**Sonila Duka, Loreta Vallja, Majlinda Vasjari, Alma Shehu, Nevila Broli**

STUDY ON TEMPERATURE CONTROL OF MASS CONCRETE 8696
**Lujun Ding, Yuhong Liu**

THE SPATIAL CHARACTERISTICS AND INFLUENCING FACTORS OF PROVINCIAL GREEN TOTAL FACTOR PRODUCTIVITY IN CHINA-BASED ON THE SPATIAL DURBIN MODEL 8705
**Xiwen Feng, Mingshang Xin, Xinghua Cui**

A NOVEL SYNTHESIS CHEMICAL CHARACTERIZATION AND BIOLOGICAL ACTIVITIES OF METAL-LEATHER PROTEIN HYDROLYSATE CHELATES 8717
**Rania Jacob, Hazem Hassan, Adel Afify, Gamal Gabr**

COMPARISON AND DISCUSSION OF THE NANHUA GLACIATION DIVISION OF THE NEOPROTEROZOIC GLACIAL PERIOD IN WESTERN TARIM BASIN, CHINA 8728
**Qilong Kang, Wanjun Wang, Zhenze Wang, Yidong Yuan**

AN EMPIRICAL ANALYSIS OF THE ENVIRONMENTAL KUZNETS CURVE IN THE HUAIHE RIVER BASIN 8740
**Jianfei Leng, Lidan Yuan**

SOIL QUALITY ASSESSMENT OF CROP ROTATIONS AND MANAGEMENT PRACTICES BASED ON MINIMUM DATA SET AND SOIL QUALITY INDICES IN THE NORTHEAST PLAIN, CHINA 8749
**Yixuan Li, Pujia Yu, Licheng Shen**

EFFECT OF SURGICAL MASK USE ON BREATHING AIR QUALITY AND OXYGEN SATURATION: A QUASI-EXPERIMENTAL MODEL 8758
**Ertan Kara, Burak Mete, Hakan Demirhindi, Ceren Kanat, Esra Mete**

EFFECTS OF HIGH VOLTAGE ELECTRIC FIELD (HVEF) TREATMENTS ON GERMINATION AND SEEDLING GROWTH OF FIELD PEA (*PISUM ARVENSE* L.) SEEDS 8763
**Taskin Erol, Kamil Kara, Ozlem Ince Yilmaz, Mustafa Dogan**

EXPERIMENTAL STUDY ON THE STRUCTURAL CONTROL OF HETEROGENEOUS DISTRIBUTION OF SHALE GAS IN EAETERN CHONGQING AREA, CHINA 8770
**Yu Zhu**

**NOTICE**

RESEARCH ON SUBSTITUTION ENERGY SUBSTITUTION IN CHINESE INDUSTRIAL SECTORS BASED ON LINEAR LOGIT MODEL 8777
**Yunsong Wang**

URBAN WATERFRONT LANDSCAPE DESIGN BASED ON ECOLOGICAL ENVIRONMENTAL PROTECTION CONCEPT AND MATERIAL APPLICATION 8786
**Fuquan Jing**

ENVIRONMENTAL INFORMATION DISCLOSURE AND HOUSEHOLD HEALTHY CONSUMPTION A STUDY BASED ON MICRO-SURVEY DATA 8793
**Yuxin Zhang, Chenghao Sun**

STUDY ON THE LAW OF GAS CAP MIGRATION OF CARBONATE CONDENSATE OIL-GAS RESERVOIR WITH WEAK VOLATILE OIL BASED ON VARIATION OF CRUDE OIL DENSITY 8802
**Chao He, Ning Wang, Qiang Zheng, Xunchen Cao, Shouheng Xiao**

© by PSP Volume 30– No. 07/2021 pages 8278-8283 Fresenius Environmental Bulletin

# LAND USE COVER CHANGE ANALYSIS (1985-2020) OF AYVALIK DISTRICT BALIKESIR TURKEY USING GEOGRAPHIC INFORMATION SYSTEMS AND REMOTE SENSING

**Figen Altiner[1,*], Abdullah Kelkit[2]**

[1]School of Graduate Studies, Department of Landscape Architecture, Canakkale Onsekiz Mart University, Canakkale, Turkey
[2]Department of Landscape Architecture, Canakkale Onsekiz Mart University, Canakkale, Turkey

## ABSTRACT

Changes in use of the urban and rural areas are increasing each passing day based on the rapid population growth and urbanization along with the developments in industry and technology. Direction and rate of this change can be detected through the calculation of the changes by the virtue of developments in technology and science. In this study, it is aimed to detect the changes which have occurred in usage of the areas between the years 1985 and 2020 in Ayvalik district, Turkey with the aid of remote sensing and geographical information systems. The land use changes calculated as the results of the study indicated that 668.6 % of the increase took place in the highway transportation network of the district, 420.7 % of which in tourism and entertainment areas, 152 % of which in the residential areas covering both the district center and villages, 133.8 % of which in the industrial areas and 32.2 % of which in the ports areas in a period of 35 years. The land use change results also showed that there has been a decrease of 19.3% in the total amount of forests that covered the landscape. Regarding the examination of the land use change on the agricultural basis, it was found out that the irrigated agricultural areas have increased by 33.4 % whereas the dry agricultural areas and the cultivated agricultural areas have decreased by %38.1 and 20.1 %, respectively.

**KEYWORDS**:
Land use change, remote sensing, geographic information systems, image classification, maximum-likelihood classification, landsat

## INTRODUCTION

Land use and change of land cover are important tools for the evaluation of different temporal and spatial changes on a global scale [1, 2, 3, 4, 5]. It has been revealed that the changes in land cover largely take place as a result of the anthropogenic activities such as deforestation, urbanization, intensification of agriculture and further destruction of land uses and etc. Particularly when the phenomenon of urbanization is considered in terms of space and time scales, it is seen that urbanization itself is the predominant anthropogenic factor which has impact on the world and its environment.[6,7,8]. The excessive population growth and the urbanization phenomenon have caused changes in urban area use and led unhealthy development of urbanization [9]. At the same time, urban land change is not only restricted to the core city, but also includes many new urban-rural spaces functionally tied to the city. The urban land change also has many impacts on rural lands and therefore deserves more attention in land-change science [10, 11, 12]

Determining the change in land use/cover in a region is the process of determining the situation of that region at different times and revealing the differences between them. To this end, accurately processed remote sensing images of different times are very important data sources [13].

Remote sensing of images has been successfully applied in many fields, such as classification process and land change detection. At the same time, the developments in satellite technologies have also made it possible to evaluate the changes that occur in urban and rural areas in the recent years. [14, 15]. Image classification is an important part of remote sensing, image analysis and pattern recognition. In some cases, the classification itself can be the subject of the analysis [16, 17]. Images received from satellites can be converted to land use and crop pattern maps by virtue of classification methods [18].

Land use/ cover maps are provided through classification of remote sensing images and they are an important part of the data which is an input to Geographic Information Systems for analysis and modeling studies [19, 20, 21]). Geographic Information Systems (GIS) is a means of an information system which provides the service of solving problem(s) by collecting data of positional units. It can also attribute data of such positional units in a portion of the earth while it can also provide their storage, mainte-

nance, analysis, evaluation, provision and acquisition of new information in this way [22, 23, 24]. The classification process in remote sensing may consist of several phases. These phases are as follows: determination of appropriate classification algorithm, data pre-processing, collecting training samples, feature extraction and post- classification processing [25].

## MATERIALS AND METHODS

Ayvalik district which is subject of this study is located within the borders of the Balikesir province on the southern coast of the Edremit Gulf and in North West part of Turkey. Location of Ayvalik district is shown in Figure 1. The district is geographically located between 39°18'54.85" and 39°17'35.17" Northern latitudes and 26° 42'52.87"-26° 45'11.38" eastern meridians [26].

The Ayvalik District which has 293 km² of acreage has a population of 70.720 according to the 2020 census data and 21.381 according to the 1985 census data issued by Turkey Statistics Institute [27] Ayvalik's winters are warm and rainy while its summers are hot and dry because of its location in the Aegean region. The dominant winds are the southwest and northeast winds. Ayvalik is one of the major tourist attraction sites of region and Turkey for its cultural and natural structure. There are many bay and headland formations in Ayvalik based on the unique feature of the Aegean region where the mountains are lined perpendicular to the sea cost. These bay and headland formations have also a positive effect on the tourism activities within the district [28].

In this study, land use changes which have occurred between the years 1985-2020 of Ayvalik district were examined. In order to examine the land use changes, remote sensing and geographical information systems were used.

Processes such as data set procurement, data pre-processing, collecting training samples, performing necessary digitization operations, classification of field uses, calculation of change and accuracy assessment were used in this study.





**FIGURE 1**
**Location of Ayvalık district**

**TABLE 1**
**Features of satellite images**

| *Satallite* | *Sensor* | *Path-Row* | *Acquisition Date* | *Number of Bands* | *Data Sources* |
|---|---|---|---|---|---|
| *Landsat 5* | *TM* | *181-033* | *21.05.1985* | *7* | *Landsat 4-5 TM C1 Level-1* |
| *Landsat 5* | *TM* | *181-033* | *19.05.1990* | *6* | *Landsat 4-5 TM C1 Level-1* |
| *Landsat 5* | *TM* | *181-033* | *17.05.1995* | *6* | *Landsat 4-5 TM C1 Level-1* |
| *Landsat 7* | *ETM* | *181-033* | *22.05.2000* | *6* | *Landsat 7 ETM + C1 Level-1* |
| *Landsat 5* | *TM* | *181-033* | *12.05.2005* | *6* | *Landsat 4-5 TM C1 Level-1* |
| *Landsat 5* | *TM* | *181-033* | *26.05.2010* | *6* | *Landsat 4-5 TM C1 Level-1* |
| *Landsat 8* | *OLI_TIRS* | *181-033* | *24.05.2015* | *8* | *Landsat 8 OLI/TIRS C1 Level-1* |
| *Landsat 8* | *OLI_TIRS* | *181-033* | *24.05.2020* | *8* | *Landsat 8 OLI/TIRS C1 Level-1* |

**Data set procurement, data pre-processing and collecting training samples.** In order to determine the land uses correctly and perform the necessary digitization processes, the following maps have been used:

Static, Stand and Function maps received from Republic of Turkey Ministry of Agriculture and Forestry,

Dams, streams and the ponds maps received from Republic of Turkey Ministry of State Water Works,

Google earth images and aerial photographs.

Landsat satellite images were utilized in order to determine the changes in the land use/cover of the Ayvalik district. A total number of 8 Landsat images belonging to the corresponding five year periods between 1985 and 2020 were used. Features of satellite images are presented in Table 1.

First, all images were checked against any defects such as striping and then all images were clipped with the purpose of covering the study area. After that, all images were corrected geometrically and radiometrically. The dark pixels in images belonging to the study area were exposed to radiometric correction using the Radiometric Correction module in the Arc Map 10.3 software program.

**Classification of land uses.** For the classification of land uses, Maximum Likelihood algorithm which is one of the Pixel-based classification methods was used in the study. Different algorithms such as Maximum Similarity, Maximum Neighborhood and etc. have been utilized in the literature for Pixel-based classification methods. Maximum Likelihood algorithm is the most commonly utilized method in pixel-based classification. Statistical values such as mean value, variance and covariance etc. are taken into account in this method. The equivalent probability curves for classes are defined and the pixels to be classified are assigned to the one with the highest probability of membership in the Maximum Similarity classification method. Furthermore, the variance-covariance matrix which will not only create brightness values of pixels, but also make differentiation for each class is formed in this method [7].

In the present study, seventeen different land cover/land use types were identified and created by using ArcMap 10.3 software. The types of land cover/land use classes are industrial areas, village centers, district centers, tourism and entertainment areas, broad-leaved forests, coniferous forests, cultivated agriculture areas, irrigated agriculture areas, dry agriculture areas, bare surfaces, lake, stream, grassland, steppe, shrubbery and meadows, highways and ports. Many representative samples were collected for each class by using area interest tool. When the signature file was created, the pixels of the image were classified into classes according to the samples by using a maximum likelihood classification. Thereby, the supervised classification command on a raster image was executed by using the signatures respectively.

**Calculation of change.** Change detection can be defined as the process of identifying differences in the state of an object or phenomenon by observing it at different times. In general, change detection involves the application of multi-temporal data sets to analyze the temporal effect quantitatively [29]. Post-classification comparison methods use separate classifications of images acquired at different times to produce difference maps. “From–to” change information can be generated. It explains to us how much change have occurred or what areas have changed during the from to period [29, 30]. In this study, post-classification change detection is employed after the eight images are classified separately.

**Accuracy Assessment.** One of the most important final steps at classification process is accuracy assessment. The aim of accuracy assessment is to quantitatively assess how effectively the pixels were sampled into the correct land cover classes. An accuracy assessment for the pixel-based land use classification was done for the images by using Arc Map 10.3 software. Reference pixels were used in order to assess the accuracy of the classification. The classified maps were statistically validated with the random validation samples collected from the Landsat satellite images. In the first phase, each and every

point had specific color tone and the pixel value which was recognized by the software itself when the data sets were trained during pixel-based land use classification. These values were considered as reference values. All the randomly generated points were then identified by the user and assigned in different classes. The correctly identified points were considered as classified values. This process was done for the 8 pixel-based classification images. In the second phase, user, producer and all accuracy matrices were run in order to assess the accuracy. User accuracy was obtained by dividing all correctly classified cells by total reference points. Producer accuracy were also obtained by dividing total correctly classified cells of land use land cover classes by total ground pixels. In order to obtain the overall accuracy, all correctly classified cells were divided by all pixels.

## RESULTS

As a result of the analysis of the changes in the land uses over the years, it has been revealed that the structural areas such as residential, industry, tourism, highways and ports have increased on a hectare basis. The classified maps of the land use of Ayvalik district are shown in Figure 2 and Figure 3.

The results of the study showed that there has been an increase of 668.6% in the highway transportation network of the district while an increase of 420.7% in tourism and entertainment areas, an increase of 152% in the residential areas in the center and villages, an increase of 133.8% in the industrial areas and an increase of 32.2% has taken place in the port areas of the district.

When the forest areas were examined, it was seen that the coniferous forest areas have increased, whereas the broad-leaved forest areas have decreased over the years. An overall 19.3% of decrease has taken place regarding the total amount of forested areas.

In line with the information obtained from the Ayvalik Forestry Directorate, it was stated that the dominant species of the area were broad-leaved forest trees until 1985. However, after 1985 broad-leaved forest trees were cut within the scope of the silvicultural implementations and the coniferous forest tree species of Pinus brutia and Pinus pinea were planted instead.





**FIGURE 2**
**Land use maps over the years (1985-1990-1995-2000) of Ayvalik district.**





**FIGURE 3**
**Land use maps over the years (2005-2010-2015-2020) of Ayvalik district.**

**TABLE 2**
**The amount of land use/cover change between 1985 and 2000**

| Land uses (Ha) | 1985 | 1990 | 1995 | 2000 |
|---|---|---|---|---|
| Industrial area | 59.88 | 71.5 | 97.84 | 107.9 |
| Village centers | 159.71 | 409.8 | 515.27 | 704.46 |
| District centers | 453.27 | 493.22 | 695.1 | 713.63 |
| Tourism and entertainment | 65.9 | 38.61 | 43.73 | 51.68 |
| Broad-leaved forest | 2295.69 | 1721.7 | 1011.22 | 715.45 |
| Coniferous forest | 2849.69 | 2610.33 | 2160.84 | 3953.91 |
| Cultivated agricultural | 10488.78 | 9762.92 | 10947.84 | 11108.77 |
| Irrigated agricultural | 1841.45 | 972.04 | 1374.22 | 1409.84 |
| Dry agricultural | 2119.19 | 2010.71 | 1383.34 | 1287.84 |
| Bare surfaces | 5812.11 | 6214.43 | 6223.66 | 5367 |
| Lake | 102.67 | 19.21 | 105.92 | 118.19 |
| Stream | 37.62 | | | |
| Grassland | 271.89 | 441.23 | 443.56 | |
| Steppe | 2021.02 | 3863.32 | 3525.74 | 3474.95 |
| Shrubbery and meadow | 616.78 | 525.8 | 648.57 | 58.52 |
| Highways | 61.58 | 122.6 | 77.94 | 180.84 |
| Ports | 52.41 | 32.22 | 54.85 | 56.66 |
| Toplam | 29309.64 | 29309.64 | 29309.64 | 29309.64 |

**TABLE 3**
**The amount of land use/cover change between 2005 and 2020**

| Land uses (Ha) | 2005 | 2010 | 2015 | 2020 |
|---|---|---|---|---|
| Industrial area | 120.66 | 117.17 | 152.57 | 140.01 |
| Village centers | 754.38 | 858.18 | 1191.12 | 1341.29 |
| District centers | 791.61 | 764.65 | 958.27 | 931.3 |
| Tourism and entertainment | 150.59 | 153.16 | 162.65 | 343.19 |
| Broad-leaved forest | 1076.43 | 1059.25 | 1201.49 | 723.28 |
| Coniferous forest | 3072.26 | 1654.6 | 2504.66 | 3915.72 |
| Cultivated agricultural | 9668.82 | 10903.71 | 10975.24 | 8384.3 |
| Irrigated agricultural | 1506.62 | 1888.6 | 2143.6 | 2457.15 |
| Dry agricultural | 1219.03 | 995.23 | 1835.61 | 1310.58 |
| Bare surfaces | 7046.11 | 7895.1 | 3502.62 | 5784.95 |
| Lake | 109.62 | 110.91 | 111.51 | 120.28 |
| Stream | 16.55 | 30.82 | 27.61 | 30.72 |
| Grassland | 109.45 | | 205.6 | 174.6 |
| Steppe | 2880.75 | 2449.86 | 3515.44 | 2884.37 |
| Shrubbery and meadow | 449.83 | 58.21 | 395.2 | 225.28 |
| Highways | 279.02 | 312.54 | 371.19 | 473.32 |
| Ports | 57.91 | 57.65 | 55.26 | 69.3 |
| Toplam | 29309.64 | 29309.64 | 29309.64 | 29309.64 |

**TABLE 4**
**Land use land cover changes of Ayvalik district during 1985–2000, 2000–2020, and 1985–2020**

| Land use | Area change (ha) | | | Percentage change (%) | | |
|---|---|---|---|---|---|---|
| | 1985-2000 | 2000-2020 | 1985-2020 | 1985-2000 | 2000-2020 | 1985-2020 |
| Industrial area | +48.02 | +32.11 | +80.13 | +80.2 | +29.7 | +133.8 |
| Village centers | +544.75 | +636.83 | +1181.58 | +341.1 | +90.3 | +739.8 |
| District centers | +260.36 | +217.67 | +478.03 | +57.4 | +30.5 | +105.4 |
| Tourism and entertainment | -14.22 | +291.61 | +277.29 | -21.5 | +564.2 | +420.7 |
| Broad-leaved forest | -1580.24 | +7.83 | -1572.41 | -68.8 | +1.1 | -68.5 |
| Coniferous forest | +1104.22 | -38.19 | +1066.03 | -38.7 | -1 | +37.4 |
| Cultivated agricultural | +619.99 | -272.47 | -2104.48 | +6 | -2.4 | -20.1 |
| Irrigated agricultural | -431.61 | +1047.31 | +615.7 | -23.4 | +74.2 | +33.4 |
| Dry agricultural | -831.35 | +22.74 | -808.61 | -39.2 | +1.7 | -38.1 |
| Bare surfaces | -445.11 | +417.95 | -27.16 | -7,6 | +7.7 | -0.5 |
| Lake | +15.52 | +2.09 | +17.61 | +15.1 | +1.7 | +17.1 |
| Stream | -37.62 | +30.72 | -6.9 | -100 | +100 | -18.3 |
| Grassland | -271.89 | +174.6 | -97.29 | -100 | +100 | -35.7 |
| Steppe | +1453.93 | -590.58 | +863.35 | -71.9 | -16.9 | +42.7 |
| Shrubbery and meadow | -558.26 | +166.76 | -391.5 | -90.5 | +284.9 | -63.4 |
| Highways | +119.26 | +292.48 | +411.74 | +193.6 | +161.7 | +668.6 |
| Ports | +4.25 | +12.64 | +16.89 | +0.08 | +22.3 | +32.2 |

The examination based on agricultural areas revealed that the irrigated agriculture areas have increased by 33.4%, whereas dry agricultural areas have decreased by 38.1% while the cultivated agricultural areas have decreased by 20.1%. The increase in irrigated agricultural activities is caused by the developments in agricultural irrigation system technologies along with the Madra dam which was built for the purpose of agricultural irrigation activities of the district. At the present time, olive groves agricultural areas constitute 80% of the total amount of the cultivated agricultural land in the district. However, population increase and the growth of the urban areas, the olive groves agricultural areas have been replaced by residential and tourism areas.

The temporal change of the land use of Ayvalik district is shown on hectare basis in Table 2 and Table 3.

The amount of land use change on hectare basis showed that there has been a decrease in grassland, shrubbery and meadow, bare surfaces and stream while there has been an increase in the steppe and lake areas (Table 4). Summary of pixel-based classification accuracy for the 8 different time frames found from accuracy assessment is shown in Tables 5–7. The highest accuracy was found for 2020 pixel-based classification (92,1% accuracy) and the lowest for 1985 (84.7% accuracy).

**TABLE 5**
**The accuracy matrix for classified images 1985, 1990,1995 and 2000**

| | 1985 | | 1990 | | 1995 | | 2000 | |
|---|---|---|---|---|---|---|---|---|
| **Classes** | Producer accuracy | User accuracy | Producer accuracy | User accuracy | Producer accuracy | User accuracy | Producer accuracy | User accuracy |
| | % | % | % | % | % | % | % | % |
| Industrial area | 75 | 80 | 81.2 | 86.6 | 81.2 | 86.6 | 81.2 | 86.6 |
| District centers | 80 | 80 | 100 | 86.6 | 100 | 86.6 | 100 | 86.6 |
| Tourism and entertainment | 73.3 | 73.3 | 80 | 80 | 92.8 | 86.6 | 75 | 80 |
| Coniferous forest | 85.7 | 80 | 65 | 86.6 | 93.3 | 93.3 | 93.3 | 93.3 |
| Cultivated agricultural | 85.7 | 80 | 92.3 | 80 | 86.6 | 86.6 | 100 | 80 |
| Irrigated agricultural | 86.6 | 86.6 | 72.2 | 86.6 | 81.2 | 86.6 | 60 | 80 |
| Dry agricultural | 92.8 | 86.6 | 92.8 | 86.6 | 76.4 | 86.6 | 80 | 80 |
| Bare surfaces | 81.2 | 86.6 | 81.2 | 86.6 | 81.2 | 86.6 | 100 | 86.6 |
| Lake | 93.3 | 93.3 | 93.7 | 100 | 93.7 | 100 | 77.7 | 93.3 |
| Village centers | 85.7 | 80 | 100 | 73 | 100 | 73.3 | 100 | 86.6 |
| Stream | 100 | 93.3 | 100 | 93.3 | 100 | 93.3 | 92.8 | 86.6 |
| Grassland | 92.8 | 86.6 | 86.6 | 86.6 | 72.2 | 86.6 | 80 | 80 |
| Steppe | 70.5 | 80 | 86.6 | 86.6 | 100 | 86.6 | 66.6 | 80 |
| Shrubbery and meadow | 81.2 | 86.6 | 100 | 86.6 | 86.6 | 86.6 | 85.7 | 80 |
| Broad-leaved forest | 100 | 100 | 86.6 | 86.6 | 87.5 | 93.3 | 100 | 86.6 |
| Highways | 86.6 | 86.6 | 80 | 80 | 85.7 | 80 | 86.6 | 86.6 |
| Ports | 80 | 80 | 100 | 86.6 | 100 | 93.3 | 100 | 86.6 |

**TABLE 6**
**The accuracy matrix for classified images 2005, 2010, 2015 and 2020**

| | 2005 | | 2010 | | 2015 | | 2020 | |
|---|---|---|---|---|---|---|---|---|
| **Classes** | Producer accuracy | User accuracy | Producer accuracy | User accuracy | Producer accuracy | User accuracy | Producer accuracy | User accuracy |
| | % | % | % | % | % | % | % | % |
| Industrial area | 76.4 | 86.6 | 100 | 93.3 | 100 | 93.3 | 82.3 | 93.3 |
| District centers | 93.3 | 93.3 | 100 | 80 | 100 | 93.3 | 100 | 100 |
| Tourism and entertainment | 100 | 80 | 88.2 | 100 | 77.7 | 93.3 | 93.3 | 93.3 |
| Coniferous forest | 100 | 86.6 | 86.6 | 86.6 | 92.8 | 86.6 | 87.5 | 93.3 |
| Cultivated agricultural | 100 | 93.3 | 100 | 86.6 | 100 | 86.6 | 87.5 | 93.3 |
| Irrigated agricultural | 62 | 86.6 | 78.9 | 100 | 73.6 | 93.3 | 100 | 93.3 |
| Dry agricultural | 85.7 | 80 | 100 | 80 | 92.3 | 80 | 100 | 86.6 |
| Bare surfaces | 76.4 | 86.6 | 86.6 | 86.6 | 100 | 86.6 | 81.2 | 86.6 |
| Lake | 87.5 | 93.3 | 93.3 | 100 | 83.3 | 93.3 | 100 | 100 |
| Village centers | 100 | 80 | 100 | 86.6 | 100 | 86.6 | 83.3 | 100 |
| Stream | 100 | 93.3 | 87.5 | 93.3 | 100 | 93.3 | 88.2 | 100 |
| Grassland | 76.5 | 86.6 | 92.8 | 86.6 | 86.6 | 86.6 | 100 | 93.3 |
| Steppe | 80 | 80 | 92.8 | 93.3 | 93.3 | 93.3 | 92.8 | 86.6 |
| Shrubbery and meadow | 85.7 | 80 | 86.6 | 86.6 | 100 | 80 | 92.3 | 80 |
| Broad-leaved forest | 85.7 | 80 | 86.6 | 86.6 | 87.5 | 93.3 | 86.6 | 86.6 |
| Highways | 81.2 | 86.6 | 92.8 | 86.6 | 83.3 | 100 | 100 | 93.3 |
| Ports | 85.7 | 80 | 92.8 | 86.6 | 87.5 | 93.3 | 100 | 86.6 |

**TABLE 7**
**Over all accuracy matrix for classified images 1985, 1990, 1995, 2000, 2005, 2010, 2015 and 2020**

| | 1985 | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2020 |
|---|---|---|---|---|---|---|---|---|
| **Over all accuracy %** | 84.7 | 85.8 | 87.8 | 84.7 | 85.5 | 89.4 | 90.2 | 92.1 |

## CONCLUSIONS

The district of Ayvalik is one of the most important tourism and agricultural locations in Turkey. As of 2017, Ayvalik has been recorded to the temporary list of UNESCO World Heritage.

The land use changes which have occurred in the district are of significant importance not only for the region but also for Turkey, in terms of economic and social development. In terms of its natural and cultural characteristics, the district has a place that should not be ignored regarding the protection and use balance. There has been a great increase in tourism, residential and industrial areas in the district, particularly after the year 2000. The increase in industrial, residential and tourism areas have caused an increase in the built-up areas and thus, urban sprawl has gradually been accelerated. The zoning for construction of many olive groves within the scope of protected areas affected agricultural activities adversely. However, the increase of the modern irrigated agricultural technologies has positively affected the general agricultural activities. The results also revealed that the forest covered areas have decreased since 1985 till the present day.

It is foreseen that this study will shed light on sustainable and future-oriented ecological, economic, social and cultural planning in this research field by determining the land use changes in Ayvalik District. The excessive population growth, urbanization, developments in industry and technology, expansion of transportation networks, natural or human activities cause disasters to change the ecosystem directly or indirectly, they have also affected and continued to affect the urban or rural areas being a part of the ecosystem.

With the increasing number of studies on determining the changes in area uses, the changes in the ecosystem can be determined over time and correct and sustainable plan decisions can be made. Studies on the determination of changes in land use can assist for making correct and sustainable plan decisions in urban and rural areas.

## ACKNOWLEDGEMENTS

This study has been prepared within the scope of the doctoral thesis named as “The examination of parameters affecting the spatial change in the Ayvalik district example”.

## REFERENCES

[1] Zhai, H., Xie, W., Li, S., Zhang, Q. (2021). Evaluation of urban ecological environment based on remote sensing based ecological index model. Fresen. Environ. Bull. 30(3), 2527-2534.

[2] Magliocca, N.R., Rudel, T.K., Verburg, P.H., McConnell, W.J., Mertz, O., Gerstner, K., Heinimann, A., Ellis, E.C. (2015). Synthesis in land change science: methodological patterns, challenges, and guidelines. Regional Environment Change. 15, 211-226.

[3] Verburg, P.H., Crossman, N., Ellis, E.C., Heinimann, A., Hostert, P., Mertz, O., Nagendra, H., Sikor, T., Erb, K.H., Golubiewski, N., Grau, R., Grove, M., Konaté, S., Meyfroidt, P., Parker, D.C., Chowdhury, R.R., Shibata, H., Thomson, A., Zhen, L. (2015). Land system science and sustainable development of the earth system: a global land project perspective. Anthropocene. 12, 29-41

[4] Li, J., Wang, Z., Lai, C., Wu, X., Zeng, Z., Chen X., Lian, Y. (2018). Response of net primary production to land use and land cover change in Mainland China since the late 1980s. Science of the Total Environ. 639, 237-247.

[5] Powers, R.P., Jets, W. (2019). Global habitat loss and extinction risk of terrestrial vertebrates under future land-use-change scenarios. Nature Climate Change. 9(4), 323-329.

[6] Patra, S., Sahoo, S., Mishra, P., Mahapatra, S.C. (2018). Impacts of urbanization on land use/cover changes and its probable implications on local climate and groundwater level. Journal of Urban Management. 7(2), 70–84.

[7] Sarvar, A., Saifulilah, S., Gulzar, S., Sohail, S., Qayyum, N., Khurshid, T. (2019). Land use land cover change analysis (2000-2018) of Tehsil Katlang, district Mardan, Pakistan, using GIS and remote sensing. Journal of Environmental & Agricultural Sciences. 20, 21-27.

[8] Noszczyk, T. (2019). A review of approaches to land use changes modeling. Human Ecologicigical Risk Assessment. 25(6), 1377-1405.

[9] Gulgun, B., Guney, M.A., Aktas, E., Yazıcı, K. (2014). Role of the landscape architecture in interdisciplinary planning of sustainable cities. Journal of Environmental Protection and Ecology. 15(4), 1877-1880.

[10] Muller, D., Munroe, D.K. (2014). Current and future challenges in land-use science. Journal of Land Use Science. 9(2), 133–142.

[11] Brenner, N., Schmid, C. (2015). Towards a new epistemology of the urban. City. 19(2), 51-182.
[12] Hersperger, A.M., Oliveira, E., Pagliarin, S., Palka, G., Verburg, P., Bolliger, J., Grădinaru, S. (2018). Urban land-use change: The role of strategic spatial planning. Global Environmental Change. 51, 32-42.
[13] Coppin, P., Lambin, E., Inge, J., Muys, B. (2002). Proceedings of the first International workshop on analysis of multi-temporal remote sensing images. University or Trento, Italy. 13-14 September 2001. eds. L. Baizzone ve P. Smits, World Scientific, N.J. 2, 455.
[14] Belgiu, M., Drăguţ, L. (2016). Random forest in remote sensing: A review of applications and future directions. ISPRS Journal of Photogrammetry and Remote Sensing. 114, 24-31
[15] Singh, P., Kikon, N., Verma, P. (2017). Impact of land use change and urbanization on urban Heat Island in Lucknow city, central India - a remote sensing based estimate. Sustainable Cities and Society. 32, 100-114.
[16] Campbell, J.B. (2002) Introduction to remote sensing, 3rd edn (New York: Guiford Press). 1-622, ISBN: 978-1-60918-176-5.
[17] Akgun, A., Eronat, A.H., Turk, N. (2004). Comparing different satellite image classification methods: an application in Ayvalik district, western Turkey. In The 4th International Congress for Photogrammetry and Remote Sensing. Istanbul. Turkey.
[18] Ozelkan, E., Saglık, A., Sumer, S.K., Bedir, M., Kelkit, A. (2018). Examination of the effect of urbanization on agricultural areas using remote sensing-a case study in Canakkale. Journal of COZUMU Faculty of Agriculture. 6(1), 123-135.
[19] Star, J., Estes, J.E. (1990). Geographic information systems: an introduction. Englewoods Cliffs. New Jersey, Prentice Hall. 1-300. ISBN:0133511235
[20] Yeh, A.G.O., Li, X. (1997). An integrated remote sensing-GIS approach in the monitoring and evaluation of rapid urban growth for sustainable development in the Pearl River Delta, China. International Planning Studies. 2(2), 193-210
[21] Ma, Z., Hart, M.M., Redmond, R.L. (2001). Mapping vegetation across large geographic areas: Integration of remote sensing and GIS to classify multisource data. Photogrammetric Engineering and Remote Sensing. 67(3), 295-307.
[22] Clarke, C.K. (1986). Advances in geographic information systems. Computers, Environments and Urban Systems. 10(3-4), 175-184.
[23] Phadke, N.D. (2006). Geographic information systems (GIS) in library and information services. The Journal of Academic Librarianship. 1(33), 152-153.
[24] Jain, M. (2009). Gis and remote sensing techniques: a case study of a developing urban center. Himansu Publications. New Delhi. 1, 1-212.
[25] Landgrebe, D.A. (2003). Signal theory methods in multispectral remote sensing. Hoboken, NJ. John Wiley and Sons. New Jersey. 1, 1-508.
[26] CTM (2020). Culture and Tourism Ministry of Turkey. From: https://balikesir.ktb.gov.tr/TR-65836/fiziki-ozellikler.html. (Accessed 05 April 2020).
[27] TSI (2020). Turkey Statistical Institute. From: http://www.tuik.gov.tr. (Accessed 10May 2020).
[28] Tibet, F.A. (2013).Analysis of Ayvalik houses by using space alignment method. Master's Thesis. Balikesir University Institude of Science.
[29] Lu, D., Mausel, P., Brondizio, E., Moran, E. (2004). Change detection techniques. International Journal of Remote Sensing. 25(12), 2365–2401.
[30] Yuan, D., Elvidge, C. (1998). NALC land cover change detection pilot study: Washington DC area experiments. Remote Sensing of Environment. 66(2), 166-178.

**Received:** **13.03.2021**
**Accepted:** **14.04.2021**

**CORRESPONDING AUTHOR**

**Figen Altiner**
School of Graduate Studies
Department of Landscape Architecture
Canakkale Onsekiz Mart University
Canakkale 17100 –Turkey

e-mail: figenaltner@yahoo.com